

Dramatic solvent effect on the volume of the Diels-Alder reaction between tetracyanoethylene and cyclopentadiene

Kiselev V., Kashaeva E., Shihab M., Medvedeva M., Kononov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The partial molar volumes (V) and the enthalpies of dissolution (Δ_{disH}) for tetracyanoethylene, cyclopentadiene, and their Diels-Alder adduct were determined at 25 °C. Eleven solvents of the π and n -donor type were used. The use of alkylbenzenes as solvents for tetracyanoethylene induces pronounced changes in the enthalpy of dissolution (up to 26 kJ mol⁻¹) and in the partial molar volume (up to 11 cm³mol⁻¹), whereas these parameters for the adduct change slightly. The V and Δ_{disH} values for cyclopentadiene virtually do not depend on the nature of the solvent. In the case of tetracyanoethylene and the adduct in n -donor solvents, considerable variations of the v and Δ_{disH} values are observed; they are not linear functions of the change in the partial molar volume of the adduct. Therefore, the reaction volumes in acetonitrile (-40.69) and ethyl acetate (-45.56) differ sharply from those in *o*-xylene (-24.28) and mesitylene (-21.76 cm³mol⁻¹).

Keywords

Diels-Alder reaction, Partial molar volume, Reaction volume, Specific interactions, Tetracyanoethylene